


Cellular and Molecular Pathophysiology of AF: *Lessons for Cardiology*

Stanley Nattel, MD
Montreal Heart Institute
University of Montreal




Mechanisms of AF: Ideas of Early 20th Century


A Ectopic focus



B Single-circuit reentry



C Multiple-circuit reentry




Engelmann, Winterberg

Lewis

Garrey, Mines


Classical Concepts: Role of Refractory Period and Wavelength Changes

A



**Wavelength (WL) =
refractory period x conduction velocity**
- minimal path length for reentry
- size of functional reentry circuits


B



**Normal atrial size
Normal WL**
- AF not sustained


**Normal atrial size
Short WL**
- AF sustained

Atrial tachycardia remodeling: AF Begets AF




Wijffels MCEF, Kirchhof CJHJ, Dorland R, et al. Circulation 1995; 92:1954-1968


Atrial tachycardia remodeling (“AF begets AF”)




Atrial tachycardia remodeling (“AF begets AF”)




Atrial tachycardia remodeling (“AF begets AF”)




Atrial tachycardia remodeling (“AF begets AF”)



Conceptual Models of AF are Linked




Single-circuit reentry




*Atrial tachycardia
remodeling*

Multiple-circuit reentry




Final common pathway

Ectopic focus




*Atrial tachycardia
remodeling*



Conceptual Models of AF are Linked






Could this process be a target
for drug therapy?



Superiority of Amiodarone in AF






Effects of antiarrhythmic drugs on AF




AT-remodeling (A+) increases AF duration.

Effects of antiarrhythmic drugs on AF






**AT-remodeling (A+) increases AF duration.
Dofetilide (DOF) and Flecainide (Flec) have no protective effects**

Effects of antiarrhythmic drugs on AF



AT-remodeling (A+) increases AF duration.
Amiodarone prevents AF-duration increase caused by AT-remodeling.

Effects of antiarrhythmic drugs on AF






AT-remodeling (A+) increases AF duration.






Amiodarone prevents AF-duration increase caused by AT-remodeling.

Is this effect of amiodarone due to remodeling prevention?





AT- remodeling suppresses α_{1c} (L-type Ca^{2+} channel) subunit expression



Amiodarone prevents AT- remodeling suppression of α_{1c} (L-type Ca^{2+} channel) subunit expression










Amiodarone prevents AT- remodeling suppression of α_{1c} (L-type Ca^{2+} channel) subunit expression




This is evidence of a direct effect to prevent remodeling

Dofetilide and flecainide do not prevent AT- remodeling suppression of α_{1c} subunit expression




Effects of agents interacting with redox mechanisms on atrial-tachycardia remodeling

Mean AF duration (D7)




AF vulnerability (D7)




Effects of agents interacting with redox mechanisms on atrial-tachycardia remodeling

Mean AF duration (D7)




AF vulnerability (D7)



Can Single-Circuit Reentry Underlie AF?


Single-circuit reentry

? →




*Atrial tachycardia
remodeling*

Multiple-circuit reentry



Final common pathway


Ectopic focus






*Atrial tachycardia
remodeling*

Electrophysiology of Sustained AF Substrate in Rapid Atrial Paced (RAP) vs CHF Dogs

RAP






CHF






*LI et al, Circulation 1999;
100:87-95*

Conduction Abnormalities and Atrial Fibrosis in CHF (Structural remodeling)





Epicardial maps



Histology




Mapping of Sustained AF in a Dog with CHF



NATTEL et al, Annu. Rev. Physiol. 2000; 62:51-77


Reentry: Stabilization by Conduction Abnormalities

A



**Wavelength (WL) =
refractory period x conduction velocity**
- minimal path length for reentry
- size of functional reentry circuits

B




**Normal atrial size
Normal WL**
- reentry unstable
- AF not sustained

**Normal atrial size
Normal WL**
Local conduction abnormality
- reentry stabilized
-Fast, irreg. condxn = AF


Conduction Abnormalities and Atrial Fibrosis in CHF (Structural remodeling)

Epicardial maps


Control






Atrial tachycardia





CHF



Could this process be a target
for drug therapy?



Can Inhibiting Ang-II Formation Prevent Development of the Substrate for AF?





E = enalapril; HI=hydralazine/isosorbide

CHF induces fibrosis and increases AF duration

Li D et al, Circulation 104: 2608-2614, 2001

Can Inhibiting Ang-II Formation Prevent Development of the Substrate for AF?





E = enalapril; HI=hydralazine/isosorbide

Enalapril attenuates fibrosis and AF duration increases

Li D et al, Circulation 104: 2608-2614, 2001

Can Inhibiting Ang-II Formation Prevent Development of the Substrate for AF?




E = enalapril; HI=hydralazine/isosorbide

Vasodilator therapy does not mimic enalapril effects


Li D et al, Circulation 104: 2608-2614, 2001

ACE antagonism and clinical AF in LV dysfunction

AF in post-MI patients with LVD





AF in SOLVD patients



Pedersen et al, Circulation 1999; 100: 376-380


Vermes et al, Circulation 2003; 107: 2926-2931

Effects of simvastatin and fenofibrate on AF duration and atrial atrial fibrosis in CHF




Factors predicting AF in patients with LV dysfunction


92% statins



Effects of fish consumption on AF occurrence





Ω-3 FA determined by ingestion



Ω-3 FA not determined by ingestion


Effects of polyunsaturated acids on structural remodeling




What is the Role of Ectopic Foci?

Ectopic focus

Single-circuit reentry




Multiple-circuit reentry




*Atrial tachycardia
remodeling*

Final common pathway



*Atrial tachycardia
remodeling*

Etopic activity from a PV



AF initiation from a PV




Other Contributors

- Atrial Ischemia
- Inflammation
- Atrial stretch


Other Contributors

- Atrial Ischemia
- Inflammation
- Atrial stretch


Occlusion of Arterial Branch to RA




Activation Mapping of AF during Atrial Ischemia



Drug effects on ischemic AF




Drug effects on ischemic AF





Other Contributors

- **Atrial Ischemia**
- **Inflammation**
- **Atrial stretch**



Effects of methylprednisolone on post-op AF in dogs









Role of Inflammation/ C-Reactive Protein



Effects of prednisone on atrial-tachycardia remodeling





Effects of prednisone on atrial-tachycardia remodeling



Effects of Glucocorticoids on AF Recurrence




Patients presenting with persistent AF randomized to receive glucocorticoid or placebo.
Active Rx= methylprednisolone, 16 mg/d x 4 wks, then 4 mg/d x 4 months.
Primary end-point = recurrence of persistent AF.





Other Contributors


- **Atrial Ischemia**
- **Inflammation**
- **Atrial stretch**


Inhibition of Atrial Fibrillation by GsMtx-4 during stretch

















CONCLUSIONS

- Atrial fibrillation can be due to one or a combination of single-circuit reentry, ectopic activity or multiple circuit reentry, with complex interactions possible
- Atrial remodeling contributes importantly to these mechanisms and may provide novel therapeutic targets
- The specific mechanisms of atrial fibrillation have important potential therapeutic consequences



감사합니다

Seoul



SN.2007

Thank you

Montreal



